

AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions, and listings, of claims in the application:


Listing of Claims:

1-23. Canceled.

24. (currently amended) A method for separating a mixture of biomolecules, comprising:

(a 1) ~~contacting the composition of claim 1~~ a composition comprising a buffer and an effective amount of a poly(M₁-g-M₂) or a salt thereof, wherein:

(a) each M₁ has the formula (I):




wherein each A₁ is independently O, S or NX₁;

each of R₁, R₂, R₃ and R₄ is independently H, C₁-C₂₀ alkyl, C₄-C₁₂ cycloalkyl, C₅-C₁₂ aryl, C₄-C₁₂ heteroaryl, -(C₁-C₂₀ alkyl)(C₅-C₁₂ aryl) or -(C₅-C₁₂ aryl)(C₁-C₂₀ alkyl);

each R₅ is independently C₁-C₂₀ alkyl, C₁-C₂₀ heteroalkyl, C₄-C₁₂ cycloalkyl, C₄-C₁₂ heterocycloalkyl, C₅-C₁₂ aryl, C₄-C₁₂ heteroaryl, -(C₁-C₂₀ alkyl)(C₄-C₁₂ cycloalkyl), -(C₄-C₁₂ cycloalkyl)(C₁-C₂₀ alkyl), -(C₁-C₂₀ heteroalkyl)(C₄-C₁₂ cycloalkyl), -(C₄-C₁₂ cycloalkyl)(C₁-C₂₀ heteroalkyl), -(C₁-C₂₀ alkyl)(C₄-C₁₂ heterocycloalkyl), -(C₄-C₁₂ heterocycloalkyl)(C₁-C₂₀ alkyl), -(C₁-C₂₀ heteroalkyl)(C₄-C₁₂ heterocycloalkyl), -(C₄-C₁₂ heterocycloalkyl)(C₁-C₂₀ heteroalkyl), -(C₁-C₂₀ alkyl)(C₅-C₁₂ aryl), -(C₅-C₁₂ aryl)(C₁-C₂₀ alkyl), -(C₁-C₂₀ heteroalkyl)(C₅-C₁₂ aryl), -(C₅-C₁₂ aryl)(C₁-C₂₀ heteroalkyl), -(C₁-C₂₀ alkyl)(C₄-C₁₂ heteroaryl), -(C₄-C₁₂ heteroaryl)(C₁-C₂₀ alkyl), -(C₁-C₂₀ heteroalkyl)(C₄-C₁₂ heteroaryl), -(C₄-C₁₂ heteroaryl)(C₁-C₂₀ heteroalkyl), -(C₁-C₄ alkyl)_qNH₂, -(C₁-C₄ alkyl)_qCONH₂, -(C₁-C₄ alkyl)NHCONH₂, -(C₁-C₄ alkyl)NHCOH or -(C₁-C₄ alkyl)_qNHCOCH₃, where each q is 0 or 1; and

each X₁ is independently H, C₁-C₂₀ alkyl, C₄-C₁₂ cycloalkyl, C₅-C₁₂ aryl, C₄-C₁₂ heteroaryl, -(C₁-C₂₀ alkyl)(C₅-C₁₂ aryl), -(C₅-C₁₂ aryl)(C₁-C₂₀ alkyl), -(C₁-C₄ alkyl)_qNH₂, -(C₁-C₄ alkyl)_qCONH₂, -(C₁-C₄ alkyl)NHCONH₂, -(C₁-C₄ alkyl)_qNHCOH or -(C₁-C₄ alkyl)_qNHCOCH₃, where each q is 0 or 1;

(b) each M₂ has the formula (II):



wherein each A_2 is independently O, S or NX_2 ;

each of R_6 , R_7 , R_8 and R_9 is independently H, C_1 - C_{20} alkyl, C_4 - C_{12} cycloalkyl, C_5 - C_{12} aryl, C_4 - C_{12} heteroaryl, $-(C_1-C_{20} \text{ alkyl})(C_5-C_{12} \text{ aryl})$ or $-(C_5-C_{12} \text{ aryl})(C_1-C_{20} \text{ alkyl})$;

each R_{10} is independently H, C_1 - C_{20} alkyl, C_1 - C_{20} heteroalkyl, C_4 - C_{12} cycloalkyl, C_4 - C_{12} heterocycloalkyl, C_5 - C_{12} aryl, C_4 - C_{12} heteroaryl, $-(C_1-C_{20} \text{ alkyl})(C_4-C_{12} \text{ cycloalkyl})$, $-(C_4-C_{12} \text{ cycloalkyl})(C_1-C_{20} \text{ alkyl})$, $-(C_1-C_{20} \text{ heteroalkyl})(C_4-C_{12} \text{ cycloalkyl})$, $-(C_4-C_{12} \text{ cycloalkyl})(C_1-C_{20} \text{ heteroalkyl})$, $-(C_1-C_{20} \text{ alkyl})(C_4-C_{12} \text{ heterocycloalkyl})$, $-(C_4-C_{12} \text{ heterocycloalkyl})(C_1-C_{20} \text{ alkyl})$, $-(C_1-C_{20} \text{ heteroalkyl})(C_4-C_{12} \text{ heterocycloalkyl})$, $-(C_4-C_{12} \text{ heterocycloalkyl})(C_1-C_{20} \text{ heteroalkyl})$, $-(C_1-C_{20} \text{ alkyl})(C_5-C_{12} \text{ aryl})$, $-(C_5-C_{12} \text{ aryl})(C_1-C_{20} \text{ alkyl})$, $-(C_1-C_{20} \text{ heteroalkyl})(C_5-C_{12} \text{ aryl})$, $-(C_5-C_{12} \text{ aryl})(C_1-C_{20} \text{ heteroalkyl})$, $-(C_1-C_{20} \text{ alkyl})(C_4-C_{12} \text{ heteroaryl})$, $-(C_4-C_{12} \text{ heteroaryl})(C_1-C_{20} \text{ alkyl})$, $-(C_1-C_{20} \text{ heteroalkyl})(C_4-C_{12} \text{ heteroaryl})$, $-(C_4-C_{12} \text{ heteroaryl})(C_1-C_{20} \text{ heteroalkyl})$, $-(C_1-C_4 \text{ alkyl})_q NH_2$, $-(C_1-C_4 \text{ alkyl})_q CONH_2$, $-(C_1-C_4 \text{ alkyl}) NHCONH_2$, $-(C_1-C_4 \text{ alkyl}) NHCOH$ or $-(C_1-C_4 \text{ alkyl})_q NHCOCH_3$, where each q is 0 or 1; and

each X_2 is independently H, C_1 - C_{20} alkyl, C_4 - C_{12} cycloalkyl, C_5 - C_{12} aryl, C_4 - C_{12} heteroaryl, $-(C_1-C_{20} \text{ alkyl})(C_5-C_{12} \text{ aryl})$, $-(C_5-C_{12} \text{ aryl})(C_1-C_{20} \text{ alkyl})$, $-(C_1-C_4 \text{ alkyl})_q NH_2$, $-(C_1-C_4 \text{ alkyl})_q CONH_2$, $-(C_1-C_4 \text{ alkyl}) NHCONH_2$, $-(C_1-C_4 \text{ alkyl})_q NHCOH$ or $-(C_1-C_4 \text{ alkyl})_q NHCOCH_3$, where each q is 0 or 1;

(c) provided that at least one M_1 is different from at least one M_2 ;

with a mixture comprising a biomolecule; and

(b 2) applying an electric field to the composition in an amount sufficient to facilitate the separation of a biomolecule from the mixture.

25. (original) The method of claim 24, wherein the separation is performed within a capillary tube and two or more biomolecules are polynucleotides.

26. (original) The method of claim 25, wherein the separation has a crossover of at least 400 base pairs.

27. Canceled.

28. (new) The method of claim 24, wherein the composition further comprises a sieve polymer.

29. (new) The method of claim 28, wherein the sieve polymer is poly(acrylamide).
30. (new) The method of claim 28, wherein the sieve polymer is poly(*N,N*-dimethyl-acrylamide) and the sieve polymer has a weight-average molecular weight of at least about 3 MDaltons.
31. (new) The method of claim 24, wherein the poly(M_1 -g- M_2) or a salt thereof has a weight-average molecular weight of from about 150,000 Daltons to about 20 MDaltons.
32. (new) The method of claim 31, wherein the composition further comprises a sieve polymer or a salt thereof having a weight-average molecular weight of from about 100,000 Daltons to about 5 MDaltons.
33. (new) The method of claim 32, wherein the sieve polymer is substantially linear poly(acrylamide).
34. (new) The method of claim 24, wherein the buffer is an aqueous buffer.
35. (new) The method of claim 34, wherein the composition has a pH of from about 5 to about 11.
36. (new) The method of claim 34, wherein the composition has a pH of from about 7 to about 10.
37. (new) The method of claim 35, wherein the composition further comprises formamide, urea, pyrrolidone, *N*-methyl pyrrolidone or a mixture thereof.
38. (new) The method of claim 35, wherein the composition further comprises urea.
39. (new) The method of claim 35, wherein the composition further comprises formamide.
40. (new) The method of claim 24, wherein M_1 is *N,N*-dimethyl-acrylamide and M_2 is acrylamide.
41. (new) The method of claim 25, wherein M_1 is *N,N*-dimethyl-acrylamide and M_2 is acrylamide.
42. (new) The method of claim 26, wherein M_1 is *N,N*-dimethyl-acrylamide and M_2 is acrylamide.
43. (new) The method of claim 28, wherein M_1 is *N,N*-dimethyl-acrylamide and M_2 is acrylamide.
44. (new) The method of claim 29, wherein M_1 is *N,N*-dimethyl-acrylamide and M_2 is acrylamide.

45. (new) The method of claim 31, wherein M_1 is *N,N*-dimethyl-acrylamide and M_2 is acrylamide.
46. (new) The method of claim 32, wherein M_1 is *N,N*-dimethyl-acrylamide and M_2 is acrylamide.
47. (new) The method of claim 33, wherein M_1 is *N,N*-dimethyl-acrylamide and M_2 is acrylamide.
48. (new) The method of claim 34, wherein M_1 is *N,N*-dimethyl-acrylamide and M_2 is acrylamide.
49. (new) The method of claim 35, wherein M_1 is *N,N*-dimethyl-acrylamide and M_2 is acrylamide.
50. (new) The method of claim 36, wherein M_1 is *N,N*-dimethyl-acrylamide and M_2 is acrylamide.
51. (new) The method of claim 37, wherein M_1 is *N,N*-dimethyl-acrylamide and M_2 is acrylamide.